

Inelastic J/ψ production at HERA in the colour singlet model with k_T -factorization

A.V. Lipatov¹

*Physical Department, M.V. Lomonosov Moscow State University,
119992 Moscow, Russia*

N.P. Zotov²

*D.V. Skobeltsyn Institute of Nuclear Physics,
M.V. Lomonosov Moscow State University,
119992 Moscow, Russia*

Abstract

In the framework of k_T -factorization QCD approach and the colour singlet model we consider J/ψ inelastic photo- and leptonproduction processes at HERA. We investigate the dependences of the single differential and double differential cross section on different forms of the unintegrated gluon distribution. The z and $\mathbf{p}_T$ dependences of the spin alignment parameter α are presented also. Our theoretical predictions agree well with recent data taken by the H1 and ZEUS collaborations at HERA. It is shown that experimental study of the polarization J/ψ mesons at low $Q^2 < 1 \text{ GeV}^2$ is an additional test of BFKL gluon dynamics.

1 Introduction

It is known that from heavy quark and quarkonium production processes one can obtain unique information on gluon structure function of the proton because of the dominance of the photon-gluon or gluon-gluon fusion subprocess in the framework of QCD [1]. Studying gluon distributions at modern collider energy (such as HERA, Tevatron) is important for prediction of heavy quark and quarkonium production cross sections at future colliders (LHC, THERA). At the energies of HERA and LEP/LHC colliders heavy quark and quarkonium production processes are so called semihard processes [2–5]. In such processes by definition the hard scattering scale $\mu \sim m_Q$ is large compare to the Λ_{QCD} parameter but on the other hand μ is much less than the total center-of-mass energy: $\Lambda_{\text{QCD}} \ll \mu \ll \sqrt{s}$. The last condition implies that the processes occur in small x region: $x \simeq m_Q/\sqrt{s} \ll 1$, and that the

¹E-mail: artem_lipatov@mail.ru

²E-mail: zotov@theory.sinp.msu.ru

cross sections of heavy quark and quarkonium production processes are determined by the behavior of gluon distributions in the small x region.

It is also known that in the small x region the standard parton model (SPM) assumptions about factorization of gluon distribution functions and subprocess cross sections are broken because the subprocess cross sections and gluon structure functions depend on a gluon transverse momentum k_T [2–5]. So calculations of heavy quark production cross sections at HERA, Tevatron, LHC and other collider conditions are necessary to carry out in the so called k_T -factorization (or semihard) QCD approach, which is more preferable for the small x region than SPM.

The k_T -factorization QCD approach is based on Balitsky, Fadin, Kuraev, Lipatov (BFKL) [6] evolution equations. The resummation of the terms $\alpha_S^n \ln^n(\mu^2/\Lambda_{\text{QCD}}^2)$, $\alpha_S^n \ln^n(\mu^2/\Lambda_{\text{QCD}}^2) \ln^n(1/x)$ and $\alpha_S^n \ln^n(1/x)$ in the k_T -factorization approach leads to the unintegrated (dependent from $\mathbf{q}_T$) gluon distribution $\Phi(x, \mathbf{q}_T^2, \mu^2)$ which determine the probability to find a gluon carrying the longitudinal momentum fraction x and transverse momentum $\mathbf{q}_T$ at probing scale μ^2 .

To calculate the cross section of a physical process the unintegrated gluon distributions have to be convoluted with off mass shell matrix elements corresponding to the relevant partonic subprocesses [2–5]. In the off mass shell matrix element the virtual gluon polarization tensor is taken in the BFKL form [2–5]:

$$L^{\mu\nu}(q) = \frac{q_T^\mu q_T^\nu}{\mathbf{q}_T^2}. \quad (1)$$

Nowadays, the significance of the k_T -factorization QCD approach becomes more and more commonly recognized [7]. It was already used for the description of a wide class heavy quark and quarkonium production processes [8–23]. It is notable that calculations in k_T -factorization approach provide results which are absent in other approaches, such as the fast growth of total cross sections in comparison with SPM, a broadening of the p_T spectra due to extra the transverse momentum of the colliding partons and other polarization properties of final particles in comparison with SPM.

We point out that heavy quark and quarkonium cross section calculations within the SPM in the fixed order of pQCD have some problems. For example, the very large discrepancy (by more than an order of magnitude) [24, 25] between the pQCD predictions for hadroproduction J/ψ and Υ mesons and experimental data at Tevatron was found. This fact has resulted in intensive theoretical investigations of such processes. In particular, it was required to use additional transition mechanism from $c\bar{c}$ -pair to the J/ψ mesons, so-called the colour octet (CO) model [26], where $c\bar{c}$ -pair is produced in the color octet state and transforms into final colour singlet (CS) state by help soft gluon radiation. The CO model was supposed to be applicable to heavy quarkonium hadro- and leptonproduction processes. However, the contributions from the CO mechanism to the J/ψ meson photoproduction contradict the experimental data at HERA for z distribution [27–30].

Another difficulty of the CO model are the J/ψ polarization properties in $p\bar{p}$ -interactions at the Tevatron. In the framework of the CO model, the J/ψ mesons should be transverse polarized at the large transverse momenta $\mathbf{p}_T$. However, this is in contradiction with the experimental data, too.

The CO model has been applied earlier [31, 32] in an analysis of the J/ψ inelastic production experimental data at HERA [33]. However, the results do not agree with each other [32].

We note that the shapes of the Q^2 , $\mathbf{p}_T^2$ and y^* distributions are not reproduced by the calculation [31], and the z distributions [32] contradict the HERA experimental data too. Results obtained within the usual collinear approach and CS model [34–37] underestimate experimental data by factor about 2.

The inelastic J/ψ production at HERA in the CS model with k_T -factorization also was considered in [15, 16]. The results [16] agree with H1 experimental data [33] both in normalization and shape only at quite small charmed quark mass $m_c = 1.4 \text{ GeV}$. The theoretical prediction [15] are stimulated the experimental analysis of J/ψ polarization properties at HERA conditions.

Recently the new experimental data on the inelastic J/ψ photo- and leptonproduction at HERA were obtained by the H1 [38, 39] and ZEUS [40] collaborations with increased statistics and precision as compared with previous experimental analyses [33]. Based on the above mentioned results here we will use the CS model and the k_T -factorization approach for the analysis of the data [38–40]. We investigate the dependences of the single differential and double differential J/ψ production cross section on different forms of the unintegrated gluon distribution. Special attention is drawn to the unintegrated gluon distributions obtained from BFKL evolution equation which has been applied earlier in our previous papers [12–16]. For studying J/ψ meson polarization properties we calculate the z and $\mathbf{p}_T$ dependences of the spin alignment parameter α .

The outline of this paper is as follows. In Section 2 we present, in analytic form, the total and differential cross section for the inelastic J/ψ photo- and leptonproduction in the CS model with k_T -factorization, and give the formulas for the relevant partonic subprocess off mass shell matrix elements. In Section 3 we present the numerical results of our calculations and compare them with the H1 [38, 39] and ZEUS [40] data. Finally, in Section 4, we give some conclusions.

2 Analytic results

In this section we calculate total and differential cross section for inelastic J/ψ photo- and leptonproduction in the CS model with k_T -factorization, and give the formulas for the relevant partonic subprocess off mass shell matrix elements.

2.1 Kinematics

As indicated in Fig. 1, we denote the 4-momenta of the incoming electron and proton and the outgoing electron, proton remnant, J/ψ meson and gluon by p_e , p_p , p'_e , p'_p , p_ψ and p_g , respectively. The initial virtual photon and BFKL gluon have a 4-momenta $q_1 = p_e - p'_e$ and $q_2 = p_p - p'_p$, so that 4-momentum transfer $Q^2 = -q_1^2$. In our analysis below we will use the Sudakov decomposition, which has the following form:

$$\begin{aligned} p_\psi &= \alpha_1 p_e + \beta_1 p_p + p_{\psi T}, & p_g &= \alpha_2 p_e + \beta_2 p_p + p_{g T}, \\ q_1 &= x_1 p_e + q_{1T}, & q_2 &= x_2 p_p + q_{2T}, \end{aligned} \quad (2)$$

where $p_{\psi T}$, $p_{g T}$, q_{1T} and q_{2T} are transverse 4-momenta of corresponding particles, and

$$p_\psi^2 = m_\psi^2, \quad p_g^2 = 0, \quad q_1^2 = q_{1T}^2, \quad q_2^2 = q_{2T}^2. \quad (3)$$

In the ep c.m. frame we can write:

$$p_e = \sqrt{s}/2 (1, 0, 0, 1), \quad p_p = \sqrt{s}/2 (1, 0, 0, -1), \quad (4)$$

where we neglect the masses of the electron and proton. The Sudakov variables are expressed as follows:

$$\begin{aligned} \alpha_1 &= \frac{m_{\psi T}}{\sqrt{s}} \exp(y_\psi), & \alpha_2 &= \frac{|\mathbf{p}_{gT}|}{\sqrt{s}} \exp(y_g), \\ \beta_1 &= \frac{m_{\psi T}}{\sqrt{s}} \exp(-y_\psi), & \beta_2 &= \frac{|\mathbf{p}_{gT}|}{\sqrt{s}} \exp(-y_g), \end{aligned} \quad (5)$$

where $m_{\psi T}^2 = m_\psi^2 + \mathbf{p}_{\psi T}^2$, y_ψ and y_g are the rapidities of J/ψ meson and final gluon respectively in the ep c.m. frame. From conservation laws we can easily obtain following conditions:

$$x_1 = \alpha_1 + \alpha_2, \quad x_2 = \beta_1 + \beta_2, \quad \mathbf{q}_{1T} + \mathbf{q}_{2T} = \mathbf{p}_{\psi T} + \mathbf{p}_{gT}. \quad (6)$$

Also the variable $z = (p_\psi \cdot p_p)/(q_1 \cdot p_p)$ is used for a description of quarkonium photo- and leptonproduction processes. In the rest frame of the proton one has $z = E_\psi/E_\gamma$.

2.2 Inelastic J/ψ leptonproduction cross section

In the k_T -factorization approach the differential cross section for inelastic J/ψ leptonproduction may be written as:

$$d\sigma(ep \rightarrow e' J/\psi X) = \frac{dx_2}{x_2} \Phi(x_2, \mathbf{q}_{2T}^2, \mu^2) \frac{d\phi_2}{2\pi} d\mathbf{q}_{2T}^2 d\hat{\sigma}(e g^* \rightarrow e' J/\psi g'), \quad (7)$$

where ϕ_2 is initial BFKL gluon azimuthal angle, $\Phi(x_2, \mathbf{q}_{2T}^2, \mu^2)$ is an unintegrated gluon distribution in the proton. The $e g^* \rightarrow e' J/\psi g'$ cross section is given by:

$$\begin{aligned} d\hat{\sigma}(e g^* \rightarrow e' J/\psi g') &= \frac{(2\pi)^4}{2x_2 s} \sum |M|_{\text{SHA}}^2(e g^* \rightarrow e' J/\psi g') \times \\ &\times \frac{d^3 p'_e}{(2\pi)^3 2p_e^0} \frac{d^3 p_\psi}{(2\pi)^3 2p_\psi^0} \frac{d^3 p_g}{(2\pi)^3 2p_g^0} \delta^{(4)}(p_e + q_2 - p'_e - p_\psi - p_g), \end{aligned} \quad (8)$$

where $\sum |M|_{\text{SHA}}^2(e g^* \rightarrow e' J/\psi g')$ is the off mass shell matrix element. In (8) $\sum$ indicates an averaging over and a sum over the final polarization states. From (7) and (8) we obtain the following formula for the inelastic J/ψ leptonproduction differential cross section in the k_T -factorization approach:

$$\begin{aligned} d\sigma(ep \rightarrow e' J/\psi X) &= \frac{1}{128\pi^3} \frac{\Phi(x_2, \mathbf{q}_{2T}^2, \mu^2)}{(x_2 s)^2 (1 - x_1)} \frac{dz}{z(1 - z)} dy_\psi \times \\ &\times \sum |M|_{\text{SHA}}^2(e g^* \rightarrow e' J/\psi g') d\mathbf{p}_{\psi T}^2 dQ^2 d\mathbf{q}_{2T}^2 \frac{d\phi_1}{2\pi} \frac{d\phi_2}{2\pi} \frac{d\phi_\psi}{2\pi}, \end{aligned} \quad (9)$$

where ϕ_1 and ϕ_ψ are azimuthal angles of the initial virtual photon and J/ψ meson respectively.

2.3 Inelastic J/ψ photoproduction cross section

As in leptonproduction case, in the k_T -factorization approach the differential cross section for inelastic J/ψ photoproduction may be written as:

$$d\sigma(\gamma p \rightarrow J/\psi X) = \frac{dx_2}{x_2} \Phi(x_2, \mathbf{q}_{2T}^2, \mu^2) \frac{d\phi_2}{2\pi} d\mathbf{q}_{2T}^2 d\hat{\sigma}(\gamma g^* \rightarrow J/\psi g'). \quad (10)$$

If we take the limit $Q^2 \rightarrow 0$ and $x_1 \rightarrow 1$, we easy obtain the following formula for the inelastic J/ψ photoproduction differential cross section in the k_T -factorization approach by analogy with the leptonproduction case:

$$d\sigma(\gamma p \rightarrow J/\psi X) = \frac{1}{16\pi (x_2 s)^2} \Phi(x_2, \mathbf{q}_{2T}^2, \mu^2) \frac{dz}{z(1-z)} \times \\ \times \sum |M|_{\text{SHA}}^2(\gamma g^* \rightarrow J/\psi g') d\mathbf{p}_{\psi T}^2 d\mathbf{q}_{2T}^2 \frac{d\phi_2}{2\pi} \frac{d\phi_\psi}{2\pi}, \quad (11)$$

We note that formulas for the the differential cross section for inelastic J/ψ photo- and leptonproduction in the usual parton model may be obtained from (9) and (11), if we take the limit $\mathbf{q}_{2T}^2 \rightarrow 0$ and average them over the transverse momentum vector $\mathbf{q}_{2T}$.

2.4 Off mass shell matrix element

There are six Feynman diagrams (Fig. 2) which describe partonic subprocess $\gamma g^* \rightarrow J/\psi g'$ at leading order in α_S and α . In the framework of the CS model and the nonrelativistic approximation the production of the J/ψ meson is considered as a production of a quark-antiquark system in the colour singlet state with orbital momentum $L = 0$ and spin momentum $S = 1$. The binding energy and relative momentum of the quarks in the J/ψ meson are neglected, resulting in $m_\psi = 2m_c$, where m_c is charm mass. The amplitude of the process $\gamma g^* \rightarrow J/\psi g'$ may be obtained from the amplitude of the process $\gamma g^* \rightarrow c\bar{c} g'$ after replacement:

$$v(p_{\bar{c}}) \bar{u}(p_c) \rightarrow \hat{J}(p_\psi) = \frac{\psi(0)}{2\sqrt{m_\psi}} \hat{\epsilon}(p_\psi) (\hat{p}_\psi + m_\psi) \frac{1}{\sqrt{3}}, \quad (12)$$

where $p_c = p_\psi/2$, $\epsilon(p_\psi)$ is a 4-vector of the J/ψ polarization, $1/\sqrt{3}$ is the color factor, $\psi(0)$ is the nonrelativistic meson wave function at the origin. The matrix element is:

$$M = e_c g^2 \epsilon_\mu(q_1) \epsilon_\sigma(q_2) \epsilon_\rho(p_g) \times \\ \times Sp \left[\hat{J}(p_\psi) \gamma^\mu \frac{\hat{p}_c - \hat{q}_1 + m_c}{(p_c - q_1)^2 - m_c^2} \gamma^\sigma \frac{-\hat{p}_c - \hat{p}_g + m_c}{(-p_c - p_g)^2 - m_c^2} \gamma^\rho \right] \quad (13)$$

+ 5 permutations of all gauge bosons. Here $\epsilon_\mu(q_1)$ and $\epsilon_\mu(q_2)$ are polarization vectors of the initial photon and gluon respectively, $\epsilon_\mu(p_g)$ is a 4-vector of the final gluon polarization. The summation on the J/ψ meson and final gluon polarizations is carried out by covariant formulas:

$$\sum \epsilon^\mu(p_\psi) \epsilon^{*\nu}(p_\psi) = -g^{\mu\nu} + \frac{p_\psi^\mu p_\psi^\nu}{m_\psi^2}, \quad (14)$$

$$\sum \epsilon^\mu(p_g) \epsilon^{*\nu}(p_g) = -g^{\mu\nu}. \quad (15)$$

The initial BFKL gluon polarization tensor is taken in form (1). For the photon we use the usual expression

$$\sum \epsilon^\mu(q_1) \epsilon^{*\nu}(q_1) = -g^{\mu\nu} \quad (16)$$

in photoproduction case and the full lepton tensor (including also the photon propagator factor and photon-lepton coupling) in leptonproduction case:

$$\sum \epsilon^\mu(q_1) \epsilon^{*\nu}(q_1) = 2 \frac{e^2}{Q^2} \left(-g^{\mu\nu} + \frac{4p_e^\mu p_e^\nu}{Q^2} \right). \quad (17)$$

For studying J/ψ polarized production we introduce the 4-vector of the longitudinal polarization $\epsilon_L^\mu(p_\psi)$ as follows [41]:

$$\epsilon_L^\mu(p_\psi) = \frac{(p_\psi \cdot p_p)}{\sqrt{(p_\psi \cdot p_p)^2 - m_\psi^2 s}} \left(\frac{p_\psi^\mu}{m_\psi} - \frac{m_\psi p_p^\nu}{(p_\psi \cdot p_p)} \right). \quad (18)$$

The evaluation of $\sum |M|_{\text{SHA}}^2$ for photo- and leptonproduction cases was done analytically by the REDUCE program. Also in our calculations we have used the JB [42] and KMS [43] parametrizations of the unintegrated gluon distributions (see also [7] for the detail information).

3 Numerical results

In this section we present the theoretical results in comparison with recent experimental data taken by the H1 [38, 39] and ZEUS [40] collaborations at HERA.

There are three parameters which determine the common normalization factor of the cross section under consideration: J/ψ meson wave function at the origin $\psi(0)$, charmed quark mass m_c and factorization scale μ . The value of the J/ψ meson wave function at the origin may be calculated in a potential model or obtained from the well known experimental decay width $\Gamma(J/\psi \rightarrow \mu^+ \mu^-)$. In our calculation we used $|\psi(0)|^2 = 0.0876 \text{ GeV}^3$ as in [44].

Concerning a charmed quark mass, the situation is not clear: on the one hand, in the nonrelativistic approximation one has $m_c = m_\psi/2 = 1.55 \text{ GeV}$, but on the other hand there are examples when smaller value of a charm mass $m_c = 1.4 \text{ GeV}$ is used [32, 45]. However, in our previous paper [16] we analyzed in detail the influence of charm quark mass on the theoretical results. We found that the main effect of change of the charm quark mass connects with final phase space of J/ψ meson, and in the subprocess matrix elements this effect is neglectable. Taking into account that the value of $m_c = 1.4 \text{ GeV}$ corresponds to the unphysical phase space of J/ψ state, in the present paper we will use value of a charm mass $m_c = 1.55 \text{ GeV}$ only.

Also the most significant theoretical uncertainties come from the choice of the factorization scale μ_F and renormalization one μ_R . One of them is related to the evolution of the gluon distributions $\Phi(x, \mathbf{q}_T^2, \mu_F^2)$, the other is responsible for strong coupling constant $\alpha_S(\mu_R^2)$. As often done in literature, we set $\mu_F = \mu_R = \mu$. In the present paper we used the following choice $\mu^2 = \mathbf{q}_{2T}^2$ as in [16, 46].

3.1 Inelastic J/ψ leptonproduction at HERA

The integration limits in (9) are taken as given by kinematical conditions of the H1 experimental data [39]. One kinematical region³ is $2 < Q^2 < 100 \text{ GeV}^2$, $50 < W < 225 \text{ GeV}$, $0.3 < z < 0.9$, $\mathbf{p}_{\psi T}^{*2} > 1 \text{ GeV}^2$ and other kinematical region is $12 < Q^2 < 100 \text{ GeV}^2$, $50 < W < 180 \text{ GeV}$, $\mathbf{p}_{\psi T}^2 > 6.4 \text{ GeV}^2$, $0.3 < z < 0.9$ and $\mathbf{p}_{\psi T}^{*2} > 1 \text{ GeV}^2$. Here and in the following, we used $\Lambda_{\text{QCD}} = 250 \text{ MeV}$.

The results of our calculations are shown in Fig. 3—5. Fig. 3 shows the single differential cross sections of the inelastic J/ψ meson leptonproduction obtained in the first kinematical region at $\sqrt{s} = 314 \text{ GeV}$. Curve 1 corresponds to the SPM calculations at the leading order approximation with the GRV (LO) gluon density, curves 2 and 3 correspond to the k_T -factorization results with the JB (at $\Delta = 0.35$ [17, 23]) and the KMS unintegrated gluon distributions. One can see that results obtained in the CS model with k_T -factorization agree very well with the H1 experimental data. The SPM calculation are lower than the data by a factor 2 — 3.

We would like to note the difference in the transverse momenta distribution shapes between curves obtained using the k_T -factorization approach and the SPM. This difference manifests the p_T broadening effect which mentioned earlier. It is visible also that only the k_T -factorization approach gives a correct description of the $\mathbf{p}_{\psi T}^2$ spectra. However, we note that the $\mathbf{p}_{\psi T}^{*2}$ distributions somewhat less well described (in contrast with $\mathbf{p}_{\psi T}^2$ spectra) at the large values of the J/ψ transverse momenta (see Fig. 3d).

Also we point out the good description of the z distributions which obtained in the k_T -factorization approach in contrast with CO model results [32], except for the region $z < 0.3$, where the contribution of the resolved photon process may be large [47].

Fig. 4 shows the single differential cross sections of the inelastic J/ψ meson production obtained in the second kinematical region at $\sqrt{s} = 314 \text{ GeV}$. Curves 1 — 3 are the same as in Fig. 3. We find also good agreement between results obtained in the CS model with k_T -factorization and H1 data. It is notable that in this kinematical region in contrast with first one the both $\mathbf{p}_{\psi T}^2$ and $\mathbf{p}_{\psi T}^{*2}$ transverse momenta distributions agree well with the experimental data.

The double differential cross sections $d\sigma/dQ^2 dz$ and $d\sigma/d\mathbf{p}_{\psi T}^{*2} dz$ (Fig. 5) obtained with k_T -factorization in the different z regions $0.3 < z < 0.6$ (Fig. 5a, b), $0.6 < z < 0.75$ (Fig. 5c, d) and $0.75 < z < 0.9$ (Fig. 5e, f) agree with the H1 data. We note that double differential cross sections $d\sigma/d\mathbf{p}_{\psi T}^{*2} dz$ somewhat less well described at the large z (see Fig. 5f). However, in this region the contribution of the diffractive processes may be large. All of these contributions are not in our consideration.

It is interesting to note that results obtained with the JB unintegrated gluon distribution at $\Delta = 0.35$ and the KMS ones, which effectively included about 70% of the full NLO corrections to the value of Δ [43], coincide practically in a wide kinematical region.

Fig. 3 — 5 show that the k_T -factorization results for inelastic J/ψ leptonproduction with realistic value of a charm mass $m_c = 1.55 \text{ GeV}$ agree well with the H1 experimental data without any additional $c\bar{c} \rightarrow J/\psi$ fragmentation mechanisms, such as the CO contributions.

3.2 Inelastic J/ψ photoproduction at HERA

The integration limits in (11) are taken as given by kinematical conditions of the H1 [38]

³Here we denote the J/ψ meson transverse momentum and rapidity in the $\gamma^* p$ c.m. frame by $\mathbf{p}_{\psi T}^*$ and y_ψ^* , respectively.

and the ZEUS [40] data. One kinematical region which corresponds to the H1 experiment is $60 < W < 240 \text{ GeV}$, $0.3 < z < 0.9$, $1 < \mathbf{p}_{\psi T}^2 < 60 \text{ GeV}^2$ and other kinematical region which corresponds to the ZEUS experiment is $50 < W < 180 \text{ GeV}$, $0.4 < z < 0.9$, $\mathbf{p}_{\psi T}^2 > 1 \text{ GeV}^2$.

The results of our calculations are shown in Fig. 6 — 8. Fig. 6 and 7 show the total and single differential cross sections of inelastic J/ψ meson photoproduction in comparison with the H1 and ZEUS data, respectively. As in previous section, curve 1 corresponds to the SPM calculations at the leading order approximation with the GRV (LO) gluon density, curves 2 and 3 correspond to the k_T -factorization results with the JB (at $\Delta = 0.35$ [17, 23]) and KMS unintegrated gluon distributions.

The W dependences of the total J/ψ photoproduction cross section at $0.3 < z < 0.9$, $0.3 < z < 0.8$ and $0.4 < z < 0.9$ are plotted in Fig. 6a, Fig. 6b and Fig. 7a respectively. One can see that results obtained in the CS model with k_T -factorization agree very well with the H1 [38] and the ZEUS [40] experimental data. The SPM results are lower than the data by a factor 2.

Concerning the shapes of the $\mathbf{p}_{\psi T}^2$ distribution (Fig. 6c and 7b), one can note a difference between the k_T -factorization and the SPM curves. As in leptonproduction case, this difference manifests the p_T broadening effect which was mentioned earlier. It is visible also that only the k_T -factorization approach gives a correct description of the H1 data.

The z distributions are shown in Fig. 6d, e, f and Fig. 7c, d, e at different $\mathbf{p}_{\psi T}$ cuts in comparison with the H1 and ZEUS data, respectively. One can see that good agreement between the k_T -factorization curves and the experimental data is observed. The z distribution somewhat are less well described at $\mathbf{p}_{\psi T} > 3 \text{ GeV}$ (see Fig. 6f). The discrepancy between the leading order SPM calculations and the experimental data is about factor 2 at $\mathbf{p}_{\psi T} > 1 \text{ GeV}$ and about order of magnitude at $\mathbf{p}_{\psi T} > 3 \text{ GeV}$ for $z \sim 0.8$. Also we note that in the region $z < 0.3$ the contribution of resolved photon process may be large [47], as in leptonproduction case.

The double differential cross sections $d\sigma/d\mathbf{p}_{\psi T}^2 dz$ (Fig. 8) in the different z regions $0.3 < z < 0.6$ (Fig. 8a), $0.6 < z < 0.75$ (Fig. 8b) and $0.75 < z < 0.9$ (Fig. 8c) are well described by the k_T -factorization approach.

It can be seen that the results obtained with the JB unintegrated gluon distribution with $\Delta = 0.35$ and the KMS ones (which effectively included the main part of the full NLO corrections to the value of Δ) practically coincide in a wide kinematical region, as in leptonproduction case.

Fig. 6 — 8 show that the k_T -factorization results for inelastic J/ψ photoproduction with realistic value of a charm mass $m_c = 1.55 \text{ GeV}$ agree well with the H1 and ZEUS experimental data without any additional $c\bar{c} \rightarrow J/\psi$ fragmentation mechanisms, such as CO contributions.

3.3 Polarization properties of the J/ψ meson at HERA

As it mentioned above, one of differences between the k_T -factorization approach and the SPM is connected with polarization properties of the final particles. In the present paper for studying J/ψ meson polarization properties we calculate the $\mathbf{p}_T$ and z dependences of the spin alignment parameter α [14–16]:

$$\alpha(\omega) = \frac{d\sigma/d\omega - 3 d\sigma_L/d\omega}{d\sigma/d\omega + d\sigma_L/d\omega}, \quad (19)$$

where σ_L is the production cross section for the longitudinally polarized J/ψ mesons, $\omega = \mathbf{p}_{\psi T}, z$. The parameter α controls the angular distribution for leptons in the decay $J/\psi \rightarrow \mu^+ \mu^-$ (in the J/ψ meson rest frame):

$$\frac{d\Gamma(J/\psi \rightarrow \mu^+ \mu^-)}{d\cos\theta} \sim 1 + \alpha \cos^2\theta. \quad (20)$$

The cases $\alpha = 1$ and $\alpha = -1$ correspond to transverse and longitudinal polarization of the J/ψ meson, respectively.

In our previous paper [16] we analyzed in detail the Q^2 and $\mathbf{p}_{\psi T}^2$ dependences of the spin parameter α in leptonproduction case. We found that it is impossible to make of exact conclusions about a BFKL gluon contribution to the polarized J/ψ production cross section because of large additional contribution from initial longitudinal polarization of virtual photons. However at low Q^2 and in photoproduction limit these contributions are negligible. This fact should result in observable spin effects of final J/ψ mesons, connected with the k_T -factorization effects. In this paper we have performed such calculations for the inelastic J/ψ photoproduction process.

The results of our calculations are shown in Fig. 9 and 10. Fig. 9 shows the parameter α as a function z and $\mathbf{p}_{\psi T}$ in comparison with the H1 experimental data which obtained in the kinematical region $60 < W < 240$ GeV, $0.3 < z < 0.9$ and $1 < \mathbf{p}_{\psi T}^2 < 60$ GeV². Curve 1 corresponds to the SPM calculations in the leading order approximation with the GRV (LO) gluon density, curve 2 corresponds to the k_T -factorization results obtained with the JB (at $\Delta = 0.35$ [17, 23]) unintegrated gluon distribution. One can see that the z dependence of the spin parameter α is not sensitive to the results of different approaches, included the nonrelativistic QCD predictions (see [38]). However, the behavior of the $\alpha(\mathbf{p}_T)$ is different in the k_T -factorization approach and the SPM (see Fig. 9b). Although the experimental points have large errors they tends to support the k_T -factorization theoretical predictions.

Fig. 10 shows the $\mathbf{p}_{\psi T}$ dependence of the spin parameter α in comparison with the ZEUS experimental data which obtained in the kinematical region $50 < W < 180$ GeV, $0.4 < z < 0.9$ (Fig. 10a and 10c) and $0.4 < z < 1$ (Fig. 10b and 10d). We note that in Fig. 10a and 10b the quantisation axis is chosen to be opposite of the incoming proton direction in the J/ψ rest frame, θ is the opening angle between the quantisation axis and the μ^+ direction of flight in the J/ψ rest frame. This frame is known as the "target frame" [40]. In Fig. 10c and 10d, the quantisation axis was defined as the J/ψ direction of flight in the ZEUS coordinate system. This frame is known as the "helicity basis" [40, 48]. Curves 1 and 2 are the same as in Fig. 9.

It is visible that only the k_T -factorization approach gives a correct description of the ZEUS data, although the experimental points have large errors. We also have large difference between predictions of the leading order of SPM and the k_T -factorization approach. The SPM predictions lies somewhat below the data at low $\mathbf{p}_{\psi T}$ and somewhat above at high $\mathbf{p}_{\psi T}$. Therefore experimental measurement of polarization properties of the J/ψ mesons will be an additional test of BFKL gluon dynamics.

4 Conclusions

In this paper we considered the inelastic J/ψ meson photo- and leptonproduction at HERA in the colour singlet model using the standard parton model in leading order in α_S and the

k_T -factorization QCD approach. We investigated the total cross section, single differential and double differential cross sections of inelastic J/ψ production on different forms of the unintegrated gluon distribution. The $\mathbf{p}_T$ and z dependences of the spin alignment parameter α presented also. We compared the theoretical results with recent experimental data taken by the H1 and ZEUS collaboration at HERA. We have found that the k_T -factorization results (in contrast with the SPM ones) with the JB and KMS unintegrated gluon distributions agree well with the experimental data at realistic value of a charm mass $m_c = 1.55 \text{ GeV}$, $|\psi(0)|^2 = 0.0876 \text{ GeV}^3$ and $\Lambda_{\text{QCD}} = 250 \text{ MeV}$ without any additional transition mechanism from $c\bar{c}$ -pair to the J/ψ mesons (such as given by the CO model). We also found that results obtained with the JB unintegrated gluon density at $\Delta = 0.35$ and KMS one, which effectively included about 70% of the full NLO corrections to the Pomeron intercept Δ , practically coincide in a wide kinematical region for J/ψ production processes at HERA conditions. Finally, it is shown that experimental study of a polarization of J/ψ meson at low $Q^2 < 1 \text{ GeV}^2$ should be additional test of BFKL gluon dynamics.

5 Acknowledgments

The authors would like to thank S. Baranov for encouraging interest and useful discussions. A.L. thanks also V. Saleev for the help on the initial stage of work. The study was supported in part by RFBR grant 02-02-17513 and INTAS grant YS 2002 N399.

References

- [1] E. Berger, D. Jones, Phys. Rev. D **23**, 1521 (1981);
S. Gershtein, A. Likhoded, S. Slabospitsky, Sov. J. Nucl. Phys. **34**, 128 (1981);
R. Baier, R. Ruckl, Nucl. Phys. B **218**, 289 (1983).
- [2] L. Gribov, E. Levin, M. Ryskin, Phys. Rep. **100**, 1 (1983).
- [3] S. Catani, M. Ciafaloni, F. Hautmann, Nucl. Phys. B **366**, 135 (1991).
- [4] J. Collins, R. Ellis, Nucl. Phys. B **360**, 3 (1991).
- [5] E. Levin, M. Ryskin, Yu. Shabelsky, A. Shuvaev, Yad. Fiz. **53**, 1059 (1991).
- [6] E. Kuraev, L. Lipatov, V. Fadin, Sov. Phys. JETP **44**, (1976) 443; **45**, (1977) 199;
Yu. Balitsky, L. Lipatov, Sov. J. Nucl. Phys. **28**, 822, (1978).
- [7] B. Andersson *et al.* (The Small x Collab.), Eur. Phys. J. C **25**, 77 (2002); hep-ph/0204115.
- [8] M. Ryskin, Yu. Shabelski, Z. Phys. C **69**, 269 (1996).
- [9] M. Ryskin, Yu. Shabelski, A. Shuvaev, Phys. Atom. Nucl. **64**, 1995 (2001); Yad.Fiz. **64**, 2080 (2001).
- [10] V. Saleev, N. Zotov, Mod. Phys. Lett. A **9**, 151 (1994).
- [11] V. Saleev, N. Zotov, Mod. Phys. Lett. A **11**, 25 (1996).

- [12] A. Lipatov, N. Zotov, Mod. Phys. Lett. A **15**, 695 (2000).
- [13] A. Lipatov, V. Saleev, N. Zotov, Mod. Phys. Lett. A **15**, 1727 (2000).
- [14] S. Baranov, A. Lipatov, N. Zotov, in *Proceedings of the 9th International Workshop on DIS and QCD (DIS'2001), Bologna, Italy, 2001*, hep-ph/0106229.
- [15] S. Baranov, Phys. Lett. B **428**, 377 (1998).
- [16] A. Lipatov, N. Zotov, hep-ph/0208237; to be published in Yad. Fiz. (2003).
- [17] S. Baranov, N. Zotov, Phys. Lett. B **458**, 389 (1999); B **491**, 111 (2000).
- [18] S. Baranov, M. Smizanska, Phys. Rev. D **62**, 014012 (2000).
- [19] P. Hagler, R. Kirschner, A. Schefer *et al.*, Phys. Rev. D **62**, 071502 (2000).
- [20] H. Jung, Phys. Rev. D **65**, 034015 (2002); hep-ph/0110034.
- [21] P. Hagler, R. Kirschner, A. Schefer *et al.*, Phys. Rev. D **63**, 077501 (2001);
F. Yuan, K.-T. Chao, Phys. Rev. D **63**, 034006 (2001); Phys. Rev. Lett. **87**, 022002 (2001), hep-ph/0009224.
- [22] H. Jung, in *Proceedings of Photon'2001, Ascona, Switzerland*, hep-ph/0110345.
- [23] S. Baranov, H. Jung, L. Jonsson *et al.*, Eur. Phys. J. C **24**, 425 (2002); hep-ph/0203025.
- [24] E. Braaten, S. Fleming, Phys. Rev. Lett. **74**, 3327 (1995);
E. Braaten, T. Yuan, Phys. Rev. D **52**, 6627 (1995).
- [25] P. Cho, A. Leibovich, Phys. Rev. D **53**, 150 (1996); D **53**, 6203 (1996).
- [26] G. Bodwin, E. Braaten, G. Lepage, Phys. Rev. D **51**, 1125 (1995); D **55**, 5853 (1997).
- [27] M. Cacciari, M. Kramer, Phys. Rev. Lett. **76**, 4128 (1996).
- [28] P. Ko, J. Lee, H. Song, Phys. Rev. D **54**, 4312 (1996); D **60**, 119902 (1999).
- [29] S. Aid *et al.* (H1 Collab.), Nucl. Phys. B **472**, 3 (1996).
- [30] J. Breitweg *et al.* (ZEUS Collab.), Z. Phys. C **96**, 599 (1997).
- [31] S. Fleming, T. Mehen, Phys. Rev. D **57**, 1846 (1998).
- [32] B. Kniehl, L. Zwirner, Nucl. Phys. B **621**, 337 (2002); hep-ph/0112199.
- [33] C. Adloff *et al.* (H1 Collab.), DESY-99-026.
- [34] J. Korner, J. Cleymans, M. Kuroda, G. Gounaris, Phys. Lett. B **114**, 195 (1982).
- [35] J.-Ph. Guillet, Z. Phys. C **39**, 75 (1988).
- [36] H. Merabet, J. Mathiot, R. Mendez-Galain, Z. Phys. C **62**, 639 (1994).
- [37] F. Yuan, K.-T. Chao, Phys. Rev. D **63**, 034017 (2001).
- [38] C. Adloff *et al.* (H1 Collab.), Eur. Phys. J. C **25**, 1, 25 (2002).
- [39] C. Adloff *et al.* (H1 Collab.), Eur. Phys. J. C **25**, 1, 41 (2002).

- [40] S. Chekanov *et al.* (ZEUS Collab.), DESY 02-163.
- [41] M. Beneke, M. Kramer, Phys. Rev. D **55**, 5269 (1997).
- [42] J. Blumlein, DESY 95-121.
- [43] J. Kwiecinski, A. Martin, A. Stasto, Phys. Rev. D **56**, 3991 (1997).
- [44] M. Kramer, Nucl. Phys. B **459**, 3 (1996).
- [45] P. Ball, M. Beneke, V. Braun, Phys. Rev. D **52**, 3929 (1995).
- [46] E. Levin, M. Ryskin, Yu. Shabelsky, A. Shuvaev, Yad. Fiz. **54**, 1420 (1991).
- [47] H. Jung, G. Schuler, J. Terron, DESY-92-028.
- [48] T. Affolder *et al.* (CDF Collab.), Phys. Rev. Lett. **85**, 2886 (2000).

Fig. 1. Diagram for $e p \rightarrow e' J/\psi X$ process.

Fig. 2. Feynmans diagram used for description partonic process $\gamma g \rightarrow J/\psi g'$ process.

Fig. 3. The single differential cross sections of the inelastic J/ψ leptonproduction obtained in the kinematical region $2 < Q^2 < 100 \text{ GeV}^2$, $50 < W < 225 \text{ GeV}$, $0.3 < z < 0.9$ and $\mathbf{p}_{\psi T}^{*2} > 1 \text{ GeV}^2$ at $\sqrt{s} = 314 \text{ GeV}$ in comparison with the H1 [39] data. Curve 1 corresponds to the SPM calculations at the leading order approximation with GRV (LO) gluon density, curves 2 and 3 correspond to the k_T -factorization QCD calculations with JB and KMS unintegrated gluon distribution.

Fig. 4. The single differential cross sections of the inelastic J/ψ leptonproduction obtained in the kinematical region $12 < Q^2 < 100 \text{ GeV}^2$, $50 < W < 225 \text{ GeV}$, $\mathbf{p}_{\psi T}^2 > 6.4 \text{ GeV}^2$, $0.3 < z < 0.9$ and $\mathbf{p}_{\psi T}^{*2} > 1 \text{ GeV}^2$ at $\sqrt{s} = 314 \text{ GeV}$ in comparison with the H1 [39] data. Curves 1 — 3 are the same as in Fig. 3.

Fig. 5. The double differential cross sections of the inelastic J/ψ leptonproduction obtained in the kinematical region $2 < Q^2 < 100 \text{ GeV}^2$, $50 < W < 225 \text{ GeV}$, $0.3 < z < 0.9$ and $\mathbf{p}_{\psi T}^{*2} > 1 \text{ GeV}^2$ at $\sqrt{s} = 314 \text{ GeV}$ in comparison with the H1 [39] data. Curves 1 — 3 are the same as in Fig. 3.

Fig. 6. The total and single differential cross sections of the inelastic J/ψ photoproduction obtained in the kinematical region $60 < W < 240 \text{ GeV}$, $1 < \mathbf{p}_{\psi T}^2 < 60 \text{ GeV}^2$, $0.3 < z < 0.9$ in comparison with the H1 [38] data. Curves 1 — 3 are the same as in Fig. 3.

Fig. 7. The total and single differential cross sections of the inelastic J/ψ photoproduction obtained in the kinematical region $50 < W < 180 \text{ GeV}$, $\mathbf{p}_{\psi T}^2 > 1 \text{ GeV}^2$, $0.4 < z < 0.9$ in comparison with the ZEUS [40] data. Curves 1 — 3 are the same as in Fig. 3.

Fig. 8. The double differential cross sections of the inelastic J/ψ photoproduction obtained in the kinematical region $60 < W < 240 \text{ GeV}$, $1 < \mathbf{p}_{\psi T}^2 < 60 \text{ GeV}^2$, $0.3 < z < 0.9$ in comparison with the H1 [38] data. Curves 1 — 3 are the same as in Fig. 3.

Fig. 9. The parameter α as a function z and $\mathbf{p}_{\psi T}$ for the inelastic J/ψ photoproduction process which obtained in the kinematical region $60 < W < 240 \text{ GeV}$, $0.3 < z < 0.9$ and $1 < \mathbf{p}_{\psi T}^2 < 60 \text{ GeV}^2$ in comparison with the H1 [38] data. Curve 1 corresponds to the SPM calculations at the leading order approximation with GRV (LO) gluon density, curve 2 corresponds to the k_T -factorization QCD calculations with JB unintegrated gluon

distribution.

Fig. 10. The parameter α as a function $\mathbf{p}_{\psi T}$ for the inelastic J/ψ photoproduction process which obtained in the kinematical region $50 < W < 180 \text{ GeV}$, $0.4 < z < 0.9$ (Fig. 10a, c), $0.4 < z < 1$ (Fig. 10b, d) and $1 < \mathbf{p}_{\psi T}^2 < 60 \text{ GeV}^2$ in comparison with the ZEUS [40] data. Curves 1 and 2 are the same as in Fig. 9.

This figure "Fig1.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig2.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig3.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig5.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig6.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig7.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig8.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig9.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>

This figure "Fig10.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/hep-ph/0210310v2>